

A service of the U.S. National Library of Medicine
and the National Institutes of Health

[\[Sign In\]](#) [\[Register\]](#)

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC
Journals Books

Search for

[Advanced Search](#)
[Text Version](#)

[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)

Entrez PubMed

[Overview](#)

[Help | FAQ](#)

[Tutorials](#)

[New/Noteworthy](#)

[E-Utilities](#)

Display Show Sort By

Send to

All: 1 Review: 0

☐ 1: [J Microencapsul. 1996 Jul-Aug;13\(4\):447-62.](#)

[Related Articles,](#)
[Links](#)

PubMed Services

[Journals Database](#)

[MeSH Database](#)

[Single Citation Matcher](#)

[Batch Citation Matcher](#)

[Clinical Queries](#)

[Special Queries](#)

[LinkOut](#)

[My NCBI](#)

Simplex lattice design for the optimization of the microencapsulation of a water soluble drug using poly(lactic acid) and poly(lactide co-glycolide) copolymer.

Elkheshen S.

Department of Pharmaceutics, Faculty of Pharmacy, Cairo University, Egypt.

Related Resources

[Order Documents](#)

[NLM Mobile](#)

[NLM Catalog](#)

[NLM Gateway](#)

[TOXNET](#)

[Consumer Health](#)

[Clinical Alerts](#)

[ClinicalTrials.gov](#)

[PubMed Central](#)

An emulsion solvent evaporation technique was applied to prepare microspheres of nicotinic acid as a water soluble vitamin. Poly(lactic acid) of two molecular weights, 2000 and 100,000 and poly(lactide-co-glycolide) (50:50) of molecular weight 18,000 were used. A simplex lattice design with the aid of a computer program was applied to study the effect of the previously mentioned polymers on different characteristics of the prepared microcapsules. The studied characteristics included the yield, the percentage drug loading and the particle size of microcapsules. Microcapsules were examined by light microscope for particle size determination and by electron scanning microscope for surface morphology characterization. The release pattern of the drug from the microcapsules was evaluated on the basis of the burst effect, the rate of release and the extent of release after 24 h. A model equation was developed to be the best representation of the relationship between the above polymers and the measured characteristics. The goodness of fit of the developed equations was checked, both statistically and experimentally. The release of the drug from microcapsules mainly followed the Higuchi diffusion model. Polylactic acid of a molecular weight 100,000 had the most pronounced delaying effect on the release rate of the drug. However, polylactic acid of a molecular weight 2,000 showed the highest improvement effect on the yield and the drug loading

characteristics of the microcapsules. Poly(lactide-co-glycolide) copolymer exhibited the lowest burst effect and the most steady drug release pattern.

PMID: 8808781 [PubMed - indexed for MEDLINE]

Display Show Sort By

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)